

2.20

(a)

Solution.

$$\int_{-\infty}^{\infty} u_0(t) \cos(t) dt = \cos 0 = 1.$$

(b)

Solution.

$$\int_0^5 \sin(2\pi t) \delta(t+3) dt = 0.$$

(c)

Solution.

$$\int_{-5}^5 u_1(1-\tau) \cos(2\pi\tau) d\tau = \int_{-5}^1 \cos(2\pi\tau) d\tau = 0.$$

2.21

(a)

Solution.

$$y[n] = \sum_{k=-\infty}^{\infty} x[n]h[n] = \sum_{k=0}^n \alpha^k \beta^{n-k} = \frac{\alpha^{n+1} - \beta^{n+1}}{\alpha - \beta}.$$

2.22

(c)


Solution. The desired convolution is

$$y(t) = \int_{-\infty}^{\infty} x(\tau)h(t-\tau) d\tau = \int_0^2 \sin(\pi\tau)h(t-\tau) d\tau.$$

This gives us

$$y(t) = \begin{cases} 0, & t < 1 \\ 2/\pi(1 - \cos(\pi(t-1))), & 1 < t < 3 \\ 2/\pi(\cos(\pi(t-3)) - 1), & 3 < t < 5 \\ 0, & t > 5 \end{cases}$$

as shown in the figure below.



2.28

(c)

Solution. Not causal because $h[n] = 2^n > 0$ for $n < 0$. Unstable because

$$\sum_{n=-\infty}^{\infty} h[n] = \sum_{n=0}^{\infty} 2^n = \infty.$$

2.29

(g)

Solution. Causal because $h(t) = 0$ for $t < 0$. Unstable because

$$\int_{-\infty}^{\infty} |h(\tau)| d\tau \geq \int_{100}^{\infty} (e^{\frac{x}{100}-1} - 1) d\tau = \infty.$$

2.33

Solution. We may solve this ODE first. Multiplying both sides by e^{2t} , we get

$$\frac{d}{dt}(e^{2t}y(t)) = e^{2t}x(t).$$

Integrating both sides from t_0 to t now gives us


$$e^{2t}y(t) - e^{2t_0}y(t_0) = \int_{t_0}^t e^{2\tau}x(\tau) d\tau.$$

For (a), let $t_0 = 0$. Then the linearity of the system follows from the linearity of the integral. For (b), since $dt = d(t - T)$, we can safely replace t with $t - T$ in the above equation without breaking its properties.

2.39

(b)

Solution. The block diagram is as shown in the figure below.




2.47

(b)

Solution. We have

$$y(t) = x(t) * h(t) = x_0(t) * h_0(t) - x_0(t-2) * h_0(t) = y_0(t) - y_0(t-2),$$

as shown in the figure below.



(d)

Solution. Not enough information.

(f)

Solution. We have

$$y(t) = \int_{-\infty}^{\infty} \frac{\partial x}{\partial t}(\tau) \frac{\partial h}{\partial t}(t-\tau) d\tau = \frac{d^2}{dt^2} \left(\int_{-\infty}^{\infty} x(\tau) h(t-\tau) d\tau \right) = \frac{d^2}{dt^2} y(t).$$

as shown in the figure below.

